Dr. Mehnaaz Ali

Research: The focus of this proposed work is to develop aptamer-based sensors for the recognition of biomarkers. Aptamers are single stranded nucleic acid sequences, which can be selected for a specific target analyte, will be used as the target recognition molecule. The proposed sensors will link the recognition of a relevant analyte by an aptamer to the activation of apoglucose oxidase via its co-factor flavin adenine dinucleotide (FAD). Specifically, this strategy will be utilized within a displacement assay where a low affinity target analog *(modified with FAD)* bound to an aptamer can be displaced by a high affinity native target. By utilizing an enzyme-based detection module with high turnover rates, this sensor aims to provide excellent signal transduction and mi

Dr. Hector Biliran

Research: *Bit1 as a tumor suppressor and a therapeutic target in NSCLC*

The work in this laboratory has made significant contributions to the field of Bit1 functional biology particularly in the context of anoikis resistance and tumorigenicity of Non-Small Cell Lung Carcinoma (NSCLC), a deadly and the most prevalent form of lung cancer. Bit1 is a mitochondrial

Immortalized HBEC has been used extensively to study the contribution of genetic alterations in the pathogenesis of lung cancer and will be used as a model system in this study. Immortalized HBEC (carrying CDK4 and hTERT) will be purchased from ATCC and cultured in K-SFM (Life Technologies, Gaithersburg, MD) medium containing 50 ug/ml bovine pituitary extract (Life Technologies) and 5 ng/ml of EGF (Life Technologies). We will stably downregulate Bit1 expression in immortalized HBECs by using lentiviral shRNA. Ready to use lentiviral Bit1 shRNA expression constructs from Open Biosystems (Huntsville, AL) have been used in our lab to successfully knockdown Bit1 expression. We will use at least two lentiviral shRNAs that target different regions of the Bit1 gene. In parallel, stable control cell line will be created via the nontargeting shRNA construct. Following confirmation of Bit1 knockdown via western blotting/real-time PCR (to be performed by undergraduate students), the stable Bit1- and control shRNA HBEC lines will be subjected to a soft agar assay to assess their anchorage-independent ability. Briefly, 1,000 viable cells will be suspended and plated in 0.37% agar in maintenance medium with or without EGF and layered over a 0.50% agar base. The results from this study will be confirmed by examining the effect of stable downregulation of AES and exogenous expression of TLE1. We have both the lentiviral AESshRNA and GFP-TLE1 expression constructs (Open Biosystems). The effect of exogenous TLE1 on the transformation ability of immortalized HBEC is particularly important in light of the recently reported lung-specific oncogenic function of TLE1.

To investigate the effect of the Bit1 pathway on the transformation potential of oncogene-induced HBECs, we will test if disruption of the Bit1 pathway will enhance transformation of Ras-induced HBECs. First, the immortalized HBEC line will be transirCab(-1(p)-10 pot)1bcb4(b)2(f)5(R)-9(L)llrge B wl0ucctv

Control and Bit1 shRNA cell lines derived from immortalized HBEC will be cultured in 3D matrigel in the presence or absence of EGF at days 5 and 8 and will be assayed for apoptosis by immunofluorescent based-TUNEL staining (Upstate Cell Signaling, Temecula, CA). The PI has extensive knowledge on TUNEL assay and will perform this work with assistance from undergraduate students. It will be interesting to see if basal apoptosis induced by 3D culture is enhanced by the lack of EGF and if this effect is attenuated by Bit1 suppression. To address if the observed apoptosis is caspase-dependent, we will prepare protein extracts from each of the cell populations by treating cultures with Dispase (Sigma) to digest the Matrigel and will be subjected to immunoblotting for inactive/cleaved caspase-3, caspase-8 and PARP (these are routine procedures to be performed by undergraduate students). To test the impact of Bit inhibition on oncogene-induced apoptosis under anchorage-independent growth condition, the control shRNA/Ras and Bit1 shRNA/Ras lines lines will be cultured in 3D Matrigel and assayed for apoptosis via TUNEL staining and detection of apoptotic markers (caspase-3, caspase-8, and PARP) by immunoblotting as described above. The specificity of Bit1 effect on basal apoptosis and oncogene-induced apoptosis in 3D cultures will be confirmed via downregulation AES and exogenous TLE1 expression in immortalized and Ras-induced HBECs.

To assess the effect of Bit1 inhibition on the invasive potential of immortalized HBEC, control shRNA and Bit1 shRNA HBEC lines will be subjected to a Matrigel/collagen 3D dimensional invasion assay as previously described. In this assay, acinar structures will be monitored for invasive behavior by measuring the development of multicellular protrusions and intercellular bridges. The use of Matrigel:collagen (1:1) mixture as opposed to pure Matrigel can enhance the formation of invasive branching morphogenesis of epithelial cells. In addition to monitoring for multicellular

factors, we will identify these DNA binding transcription factor(s) via a large scale co-immunoprecipitation. Here, TLE1 will be immunoprecipitated from controlshRNA/TLE1shRNA and control/GFP-TLE1 cells cultured in attached or detached conditions. The PI has extensive experience and is knowledgeable of the constraints associated with this assay. We have specific anti-TLE1 (Abcam) as well as anti-GFP (Sigma) antibodies that are functional for immunoprecipitation. Because of close homology of TLE1 with other TLE homologues (such as TLE2 and TLE3), we will deplete the TLE homologues (using TLE2 and TLE3 specific antibody) in the sample prior to the actual specific TLE1 immunoprecipitation. The resulting specific co-precipitating proteins will be identified using mass spectroscopy (LCRC proteomics facility). Proteomic data will not only be validated by co-immunoprecipitation studies with specific antibodies but will be analyzed for DNA binding/transcriptional activity and their association with cell attachment or detachment. Importantly, the promoter of promising TLE1 target gene(s) derived from microarray studies will be examined for

observed when planktonic bacteria derived from the same cultures were not physically associated with the solid pyrite. The premise is that these electron dense bodies repre- sent nanoparticles of insoluble substrate that have been cleaved from the exposed surface of the solid and distributed around the outer periphery of the bacterium.

Student Project #3

Transmission Electron Microscopic Analyses of Bacteria that Create Corrosion Tunnels in Solid Pyrite

The hypothesis of this student project is that acidophilic bacteria that respire aerobically on insoluble pyrite create corrosion tunnels within the solid mineral that are approximately the dimensions of individual bacteria. We formulated this hypothesis based on microscopic observations obtained when *Acidithiobacillus ferrooxidans* respired at the expense of particulate pyrite (1-3), as well as our own observations that aerobic rer1 3j(a)0.3 0.0 7 1 Tc(-0.0 8 2 Tw 3 0 0 Td (0 1 J d 0 (e 0 j w 8 (

Dr. Jeremy Cohen

Research: In the Fall 2012 semester, Dr. Jeremy Cohen began a tenure track Assistant Professorship in the Department of Psychology at Xavier University of Louisiana. Dr. Cohen is a Cognitive Neuroscientist with an undergraduate degree in Cell and Molecular Biology from Tulane University and a PhD in Neuroscience from Tulane University. His graduate work focused on studying human brain anatomy related to persistent developmental stuttering. From there his research interests0 5Fd (y)Tj 0.480 Tw ( )Tj (br)

making enabled by polymerization reaction monitoring and control technologies could thereby provide significant returns across all sectors, and represents a strategic opportunity for research investment that will sustain and enhance regional competiveness into the future.

II. Synthesizing Dendrimer with Precisely Defined Numbers of Click Ligands for Target Delivery Application

Nanomaterials provide a potential platform for cancer treatment due to their ability to simultaneously detect early stage cancers, target cancer cells over non-cancerous cells, directly deliver an active anti-cancer therapeutic, and monitor treatment efficacy. Poly(amidoamine) (PAMAM) dendrimers are a class of nanomaterials ideally suited for use as a drug delivery platform due to their controllable size, uniformity, and ample number of versatile terminal amine groups that allow for effective conjugation reactions. Thus, functionalizing dendritic polymers with small-molecule conjugates yields nanomtgl

Tj 0.28 jfl)Tj 31 0 T j[E[(rl)]4.00253(l >2(a)Tj 01.41 36n.7541 0 T jfl)Tj 3547h e7(i3t)Tj 1.10o0 T13080 0 l)]kTd [

Dr. Maryam Foroozesh

Cytochrome P450 enzymes are the most important enzymes responsible for the biotransformation and metabolism of endogenous and xenobiotic compounds including procarcinogens and pharmaceuticals. The long-term goals of many of our projects are to design mechanism-based irreversible inactivators selective for human cytochrome P450s 1A1, 1A2, 2A6 and 2A13. Selective mechanism-based (suicide) inhibitors are valuable experimental tools for studying the active sites of P450 enzymes.
Over the past years, our effort

We will explore possibilities for the student to receive an off-campus summer experience, however ex

will be tested using nuclear extracts from several different cell lines, including those that do not express PAX3-FOXO1. The student will use Western Blot to identify proteins that associate with PAX3-FOXO1 in the DNA complex. If multiple proteins are isolated, MALDI TOF MS will be used to determine the identities of the interacting proteins Immunoprecipitations using PAX3- FOXO1 as bait will also be performed under the same conditions to determine if the interaction between PAX3-FOXO1 and other proteins requires4 0 S5a

Student Project: Overall goal of the student project is to engage students in the research activities in the area of the dynamics and asymptotic behavior of discrete systems. Measurable outcomes are student's presentations (poster and oral) at students' meetings and at Joint Mathematics Meetings (annual joint meeting of all mathematical societies and associations), preparation of the joi

Dr. Kathleen Morgan

Current research in the Morgan lab focuses on simple carbohydrates and related compounds such as diols. Carbohydrates are ubiquitous in nature, yet there are still major gaps in our knowledge of sugar thermochemistry. Heats of formation ($_f$) are being determined for simple monosaccharides having 2-4 carbons; remarkably, these fundamental values are unknown. A combination of reaction calorimetry, vaporization measurements, and high-level calculations will be used to obtain the $_f$ for sugars and related compounds such as 1,2-diols. Although high-level calculations can often be used to obtain high quality thermochemical data, the compounds of interest have a large number of important conformations, making accurate calculations far from routine. The $_f$ determined in this study will be used in conjunction with bond energies determined by collaborators, which afford the $_f$ of free radicals that result from the degradation of carbohydrates.

II.

Thermochemical studies of diols: A student working on this project would start by determining reaction conditions for a reaction calorimetry study of diols. One reaction that seems promising is the hydrolysis of epoxides, and the student would need to further optimize the reaction conditions for our purposes. Once complete, this reaction would be used to measure the heats of reaction for a series of diols, via reaction calorimetry. At the same time, the student can work on calculations of the target diols to include various conformations. Other experiments are possible, such as determination of heats of vaporization and/or sublimation.

Thermochemical studies of simple sugars: A student working on this project would verify that the reaction conditions that have been developed will work on the carbohydrates planned for study, which will include glyceral and dihydroxyacetone. Then, the student will help with reaction calorimetry experiments that will, eventually, afford the heats of formation for the target compounds. The student will also work on calculations of the sugars and their reduced forms, in various conformations. Other experiments are possible, such as determination of heats of vaporization and/or sublimation.

Dr. Harris McFerrin

Although always focusing on the development and regulation of blood vessel formation, my research over the last ten years has touched on a number of topics: Human Immunodeficiency Virus (HIV), Human Herpes Virus-8 (HHV8), Herpes simplex virus-1 (HSV-1), Tumor Angiogenesis, Diabetes and Adult Human Mesenchymal Stem Cells (hMSC). These topics are all linked by their interaction with the vasculature through the up-regulation or down-regulation of the survival and growth of blood vessels. For example. HIV and HHV8 increase survival and growth of blood vessels to form lesions commonly known as Kaposi sarcoma. Diabetes negatively affects the vasculature, causing decreased blood flow and is linked to atherosclerosis and diabetic retinopathy. My lab has shown that the adult stem cells known as hMSC help to form blood vessels *in vivo* and are able to roll on and adhere to the vasculature through specific interactions, suggesting that they may be able to "home" to sites of damage.

Currently, my laboratory is investigating the role of cyclin-dependent kinases in HSV-1

infection of the eye. HSV-1 infects greater than 90% of humans worldwide and during ocular infection produces inflammation and neovascularization that can lead to blindness. In the United States, HSV infection is the leading cause of infection-induced blindness; nearly 40,000 new cases are reported, and 300,000 cases are treated yearly. We have demonstrated in a mouse eye model of HSV-1 infection that flavopiridol (a cyclin-depend kinase inhibitor) reduces neovascularization of the cornea as well as trifluorothymidine, the gold standard of treatment, and an international patent was submitted based on this research.

Dr. Krista Mincey

My research is looking at prostate cancer perception and risk in young Black males. Through focus groups with college-aged Black males 18-29, I seek to understand their prostate cancer knowledge and what behaviors or elements put them at greater risk for prostate cancer in the future. This information can then be used to inform prostate cancer programs geared toward prevention in young Black men

Dr. Patience Obih

Title: ***Preparation and Screening of Medicinal Herbs for Antidiabetic Activity.***

Research

There are about 24 million Americans living with diabetes. Every year, 1.3 million are diagnosed with type 2 diabetes. Diabetes mellitus is a chronic disease associated with many debilitating complications which result in substantial morbidity and mortality. Though medicinal plants have been used historically throughout the world, few of them have been validated scientifically. Many studies have been able to isolate the compounds but have not gone beyond that to test them on animal or in human. The World Health Organization (WHO) is in support of the use of herbal remedies for the management of diabetes. Medicinal herbs are less likely to have drawbacks of the conventional drugs used for diabetes. Further research is required to validate the antidiabetic effects of medicinal plants. As a result of the limitations in many existing conventional antidiabetic drugs, it is crucial to search for alternative remedies for diabetes mellitus, and this has led many investigators to investigate traditional remedies for better alternatives. These traditional medicines are hoped to be cheaper, efficacious, more tolerable, and produce less side effects. The aim of our research therefore is to prepare and screen medicinal extracts for their antidiabetic activities. Our first goal is to examine the effects of medicinal herbs, Bitter leaf (*Vernonia amygdalina*), Dandelion (T*araxacum officinale*), Bitter melon (*Mormodica charantia),* Nchanwu (*Orignum vulgare,* Bitter Kola (*Garcinia kola*) on hyperglycemia using a glucose tolerance test on streptozotocin-induced diabetic mice. The second goal is to examine the mechanism of action of the medicinal herbs by evaluating their ability to inhibit alpha glucosidase and lipase enzymes.

Student Involvement:

In year 1(t)-1 Tc Tf -0.013-10h77h/a

goal 1.

In yea

1. ***To demonstrate that nutritionally deprived noncancerous MCF-10A cells undergo quiescence.*** Based on our preliminary data, our working hypothesis is that nutritionally-deprived MCF-10A cells undergo senescence, thereby promoting metastasis of TNBC cells. Our gene expression profile indicated that TNFSF10, or TRAIL is highly expressed in MCF-10A cells starved verses non-starved MCF-10A cells, suggesting these starved, noncancerous cells may undergo apoptosis and

Florastina Payton-Stewart

Resear

Jayalakshmi Sridhar

Research: The objective of the proposed research is to find inhibitors of HER2 oncogenic isoform HER2_16 that can inhibit the growth drug-resistant breast cancer. The long term goals of this project are to provide therapeutic agents for the treatment of aggressive and drug-resistant breast cancers. [(s)im.34 0 0 Tw 0.95 0 Td (re)Tj 0 Tc 0 Tw 0.78 0 Td (aj 0 Tc 0 T -0.004 Tc 0.004 8rh[4

Zhe Wang

Research: Controlling the film forming parameters is the key challenge in this regard, having implications on sensitivity, stability and reproducibility, interference and coupling, and electrochemical connectivity factors of the sensor performance. In the course of this project, inverted mesa structures will be designed, fabricated, and validated to address these challenges of E-QCM integration on monolithic QCM wafers. Finalized

The student will have completed the